

# Effects of Long-term Tillage and Crop Rotation on Soil Carbon and Soil Productivity

Mahdi Al-Kaisi, assistant professor  
Mark Licht, program specialist  
Department of Agronomy

## Introduction

Tillage system and crop rotation significantly affect long-term soil productivity and soil quality components such as soil carbon and other soil physical and chemical properties. In addition, both tillage and crop rotation affect weed and soil disease control. There is a definite need for well-defined, long-term tillage and crop rotation studies across the different soil and climatic conditions in the state. The objective of this study is to evaluate the long-term wide range of effects of different tillage systems and crop rotations on soil productivity.

## Materials and Methods

This study was conducted on eight Iowa State University research and demonstration farms in 2002. Treatments included five tillage systems (no-till, strip-tillage, chisel plow, deep ripper, and moldboard plow) and two crop rotations of corn-corn-soybean and corn-soybean across the five tillage systems and several soil associations. Initial soil samples were collected from sites that were ready to implement tillage treatments during the spring of 2002. Sites that did not implement tillage treatments in spring of 2002 were planted to bulk corn and were soil sampled in fall 2002 prior to implementing tillage treatments. The soil samples, collected from all sites for depths 0–6, 6–12, 12–18, and 18–24 inches, will be analyzed for total carbon and total nitrogen. The experimental design was a randomized complete block design with four replications. Plot size ranged from 30–50 ft in width (depending on the location) and was approximately 90 ft long.

Yield will be determined from the center rows (number of rows will vary by location). The long-term effects of tillage and crop rotation on

total soil carbon and total nitrogen will be monitored biyearly, or more frequently. Seasonal measurements such as nitrogen use efficiency, soil bulk density, infiltration rate, etc., may be conducted on selected sites (depending on availability of funding).

## Results and Discussion

Results of first year (2002) yields of corn and soybeans for the Southeast Research Farm are summarized in Figures 1 and 2. The corn yields of no-till and strip-tillage were significantly less than the other three tillage systems (chisel plow, moldboard plow, and deep rip). No-till and strip-tillage yields were 13 bushels/acre less than the moldboard plow, deep ripper, and chisel plow tillage systems. In general, corn yields ranged from 194.5 bushels/acre to 179 bushels/acre.

Soybean yields of all tillage systems were not significantly different. Soybean yields ranged from 52 bushels/acre to 49 bushels/acre. First year tillage systems do not seem to make a significant difference in corn or soybean yields.

## Acknowledgments

We would like to thank Kevin Van Dee and Matt Hunt for their time and labor for plot setup, planting, and harvesting.




Figure 1. Effect of tillage system on corn yield in 2002 at Crawfordsville, IA.




Figure 2. Effect of tillage system on soybean yield in 2002 at Crawfordsville, IA.